


Routing entanglement through quantum networks

Karl Pelka^{✉,*}, Matteo Aquilina[✉], and André Xuereb^{✉,†}

Department of Physics, University of Malta, Msida MSD 2080, Malta

 (Received 14 February 2024; revised 13 November 2025; accepted 20 April 2026; published 29 June 2026)

Entanglement, one of the clearest manifestations of nonclassical physics, holds significant promise for technological applications such as more secure communications and faster computations. In this Letter, we explore the use of nonreciprocal transport in a network of continuous-variable systems to route entanglement in one direction through the network. We develop the theory and discuss a potential realization of controllable flow of entanglement in quantum systems; our method employs only Gaussian interactions and engineered dissipation to break the symmetry. We also explore the conditions under which thermal fluctuations limit the distance over which the entanglement propagates and observe a counterintuitive behavior between this distance, the strength of the entanglement source, and the strength of the hopping through the network.

DOI: [10.1103/hsc8-pv2s](https://doi.org/10.1103/hsc8-pv2s)

Introduction. Applications of quantum technologies [1] rely on the exploitation of effects such as interference and entanglement [2] that are hallmarks of quantum mechanics. The potential that lies in entangled systems—correlations that surpass, and cannot be explained, by classical physics—is so promising that studies of networks [3] envisioning a quantum internet [4] have created entire research fields [5–7]. A useful resource for distributing entanglement in distant nodes is found in propagating photons [8], enabling quantum key distribution through space using entanglement-based schemes [9]. Here, we explore the propagation of entanglement in a cascaded quantum system, showing that it is, in principle, possible to set up a scheme that allows entanglement to percolate one way, but not the other, through a network of quantum devices. Our exploration is phrased in the language of optomechanics and phonons, but can be applied to any set of interacting continuous-variable quantum systems.

The significance of nonreciprocity in quantum information applications is established in a variety of architectures, ranging from photonic setups [10–12] and superconducting circuits [13,14] to optomechanical systems [15–20]. In a largely separate line of research, the dynamics of micromechanical systems subject to radiation pressure forces has been explored thoroughly within the field of optomechanics [21,22]. This optomechanical interaction has also been used to explore the many-body dynamics of systems of coupled optomechanical networks, giving rise to phenomena including the possibility of obtaining stronger coupling at the single-photon level [23,24], topological physics [25–27], dynamical

gauge fields [28], and synchronization [29–37]. Seminal work has shown the cooling of motion of the mechanical system down to its ground state [38,39] and generation of quantum entanglement between mechanical oscillators [40,41]. A recent focal point of interest within optomechanics lies in the research on nonreciprocal manipulation of fluctuations in quantum systems, including routing thermal noise with optomechanical thermal rectifiers [42] and swapping entanglement in hot modes using cold auxiliary modes [43].

In this Letter, we employ the vision of a quantum-communication network that consists of a number of nodes in a given geometry sharing some quantum correlations described by a quantum state ρ . We demonstrate that along a connected quantum channel a combination of coherent hopping and engineered dissipation enables or disables the flow of entanglement to another node in accordance with the implemented nonreciprocity, effectively achieving an entanglement routing device. Our results show that the creation of entanglement through squeezing with our scheme increases the effective temperature in the channel and therefore limits the propagation of entanglement through the system by impairing the system's stability. Nevertheless, we find that entanglement can indeed propagate throughout a channel of several nodes and that state-of-the-art optomechanical experiments provide a basis that could realize our scheme.

We proceed by first deriving the effective quantum optics model based on cascaded quantum systems. This model will give us the physical intuition for the effects that occur and their limitations. Subsequently, we employ an already established mapping to an optomechanical model where the mechanical degrees will replace the bath degrees and show that the mechanism can be realized in experiments with achievable parameters. Ultimately, we finish with a short outline on the implications of our results.

Theoretical model. To start off, we analyze the collective dynamics of a chain of M bosonic modes acting as nodes in a transport channel with $M - 1$ nonreciprocal edges [10,42] coupled to a squeezed bosonic mode described by the

*Contact author: karl.a.pelka@um.edu.mt

†Contact author: andre.xuereb@um.edu.mt

Hamiltonian

$$\hat{H}/\hbar = \sum_{k=0}^M [\omega_k \hat{a}_k^\dagger \hat{a}_k + (iJ_k \hat{a}_k^\dagger \hat{a}_{k+1} + \text{H.c.})] + \frac{ir}{2} (\hat{a}_0^2 - \hat{a}_0^{\dagger 2}), \quad (1)$$

with \hat{a}_k describing the bosonic annihilation operators of the modes, ω_k their respective resonance frequencies, J_k the hopping rates between neighboring nodes, and r the squeezing parameter of the squeezed bosonic mode $\hat{a}_0 \equiv \hat{a}_{M+1}$. We impose cascaded quantum dynamics (see Sec. 12.1 of Ref. [44]) onto the transport channel by including common heat baths connecting neighboring oscillators \hat{a}_l and \hat{a}_{l+1} for $l \in (1, \dots, M-1)$. The oscillators decay with the rate γ into their common bath with average occupation $\bar{N}_{l,l+1}$, realized with the standard dissipator in Lindblad form $\mathcal{D}_\delta[\rho] = \delta\rho\delta^\dagger - \frac{1}{2}\{\rho, \delta^\dagger\delta\}$ through the collective operators $\hat{c}_l = (\hat{a}_l + \hat{a}_{l+1})/\sqrt{2}$ obeying the bosonic commutation relation $[\hat{c}_l, \hat{c}_l^\dagger] = 1$. This turns each of the $M-1$ edges between oscillator \hat{a}_l and \hat{a}_{l+1} into a unidirectional transport edge through the matching condition $J_l = \gamma/2$ [10,42]. The dynamics of the system's density matrix ρ are described by the master equation in Lindblad form $\dot{\rho} = -(i/\hbar)[\hat{H}, \rho] + \sum_{l=1}^{M-1} \gamma \{(\bar{N}_{l,l+1} + 1)\mathcal{D}_{\hat{c}_l}[\rho] + \bar{N}_{l,l+1}\mathcal{D}_{\hat{c}_l^\dagger}[\rho]\}$. The physical interpretation of the nonreciprocal effect is based on the interference of the direct hopping term from an oscillator to the next and the indirect path of hopping to the same oscillator through the common bath [10,42]. The nonreciprocity relies on the coherent addition of these two paths in the direction $k \rightarrow k+1$, whereas the two terms cancel each other in the opposite direction $k \rightarrow k-1$ (see Sec. 12.1 of Ref. [44]). To allow for more generality, we admit decay with the rate γ_{out} into a distinct bath with average occupation \bar{N}_k for each of the oscillators \hat{a}_k and obtain the master equation

$$\dot{\rho} = -\frac{i}{\hbar}[\hat{H}, \rho] + \sum_{k=0}^M \gamma_{\text{out}} \{(\bar{N}_k + 1)\mathcal{D}_{\hat{a}_k}[\rho] + \bar{N}_k \mathcal{D}_{\hat{a}_k^\dagger}[\rho]\} + \sum_{j=1}^{M-1} \gamma \{(\bar{N}_{j,j+1} + 1)\mathcal{D}_{\hat{c}_j}[\rho] + \bar{N}_{j,j+1} \mathcal{D}_{\hat{c}_j^\dagger}[\rho]\}. \quad (2)$$

Figure 1 depicts a schematic illustration of the system of interest. The first panel, Fig. 1(a), shows a small-scale model consisting of a squeezed mode \hat{a}_0 and three modes \hat{a}_1 to \hat{a}_3 that constitute a prototypical unidirectional chain. These modes are coupled directly with their nearest neighbors through hopping terms with rate J_k in the Hamiltonian in Eq. (1). Each of the modes is coupled to a bath into which its excitations can decay at a rate γ_{out} . In addition, the nodes that constitute the unidirectional chain are pairwise connected through a common bath into which the connected modes can decay at the rate γ . Figure 1(b) presents an abstract representation of this model by reducing nodes to vertices and showing directionality between vertices by arrows. Figure 1(c) generalizes this model to any number M of modes in the chain, whereas Fig. 1(d) effectively reverses the direction of propagation in the chain by connecting the squeezed mode to the opposite end of the chain.

We proceed by converting the master equation from Eq. (2) into equivalent quantum Langevin equations.




FIG. 1. Schematic illustrations of a squeezed oscillator connected to chains consisting of unidirectionally coupled nodes. (a) The squeezed node \hat{a}_0 is connected directly to one end of the chain constituting the quantum channel and the constituents of the channel are connected to their neighbors directly with rates J_k as well as indirectly through a common bath in which the oscillators can decay excitations at rates γ . All nodes can decay excitations into distinct baths at rates γ_{out} . (b) Abstract depiction of this chain and the directionality of the connections in this system. (c) Abstraction of the node \hat{a}_0 connected to a quantum channel of M nodes in the forward direction. (d) Depiction of the node \hat{a}_0 connected to a quantum channel of M nodes in the backward direction.

Therefore, we derive the operator equations and use the fluctuation-dissipation theorem to add noise operators for the distinct baths \hat{b}_k^{in} that obey $\langle (\hat{b}_k^{\text{in}})^\dagger(t) \hat{b}_m^{\text{in}}(t') \rangle = \bar{N}_k \delta_{km} \delta(t-t')$, $\langle \hat{b}_k^{\text{in}}(t) (\hat{b}_m^{\text{in}})^\dagger(t') \rangle = (\bar{N}_k + 1) \delta_{km} \delta(t-t')$ and for the common baths $\hat{b}_{l,l+1}^{\text{in}}$ obeying $\langle (\hat{b}_{l,l+1}^{\text{in}})^\dagger(t) \hat{b}_{m,m+1}^{\text{in}}(t') \rangle = \bar{N}_{l,l+1} \delta_{lm} \delta(t-t')$, $\langle \hat{b}_{l,l+1}^{\text{in}}(t) (\hat{b}_{m,m+1}^{\text{in}})^\dagger(t') \rangle = (\bar{N}_{l,l+1} + 1) \delta_{lm} \delta(t-t')$, while all other combinations vanish under the white-noise assumption. The resulting equations for the corresponding quadrature operators $\hat{x}_\delta = (\hat{\phi} + \hat{\phi}^\dagger)/\sqrt{2}$ and $\hat{p}_\delta = -i(\hat{\phi} - \hat{\phi}^\dagger)/\sqrt{2}$ can be summarized as $\dot{\vec{r}} = \mathcal{A}\vec{r} + \mathcal{B}\vec{r}_{\text{in}}$ for the quadrature vector $\vec{r} = (\hat{x}_{\hat{a}_0}, \hat{p}_{\hat{a}_0}, \dots, \hat{x}_{\hat{a}_M}, \hat{p}_{\hat{a}_M})^T$ and noise quadrature vector $\vec{r}_{\text{in}} = (\hat{x}_{\hat{b}_0^{\text{in}}}, \hat{p}_{\hat{b}_0^{\text{in}}}, \dots, \hat{x}_{\hat{b}_M^{\text{in}}}, \hat{p}_{\hat{b}_M^{\text{in}}}, \hat{x}_{\hat{b}_{1,2}^{\text{in}}}, \hat{p}_{\hat{b}_{1,2}^{\text{in}}}, \dots, \hat{x}_{\hat{b}_{M-1,M}^{\text{in}}}, \hat{p}_{\hat{b}_{M-1,M}^{\text{in}}})^T$ in terms of a dynamical matrix $\mathcal{A} \in \mathbb{R}^{2(M+1) \times 2(M+1)}$ and an adjacency matrix $\mathcal{B} \in \mathbb{R}^{2(M+1) \times 4M}$, which are documented in the Supplemental Material [45]. Since the equations are linear, the state of the system at any point in time is fully described by the first $\langle \vec{r}_j \rangle$ and second moments of the quadrature operators. The second moments can be arranged into a covariance matrix $\mathcal{V}_{ij} = \langle \vec{r}_i \vec{r}_j + \vec{r}_j \vec{r}_i \rangle - 2\langle \vec{r}_i \rangle \langle \vec{r}_j \rangle$. Its dynamics can be shown to obey the Lyapunov equation $\dot{\mathcal{V}} = \mathcal{V}\mathcal{A}^T + \mathcal{A}\mathcal{V} + \mathcal{N}$, with the noise matrix \mathcal{N} (see the Supplemental Material [45]) having elements $\mathcal{N}_{ij} = \sum_q (\bar{N}_q + \frac{1}{2})(\mathcal{B}_{iq} \mathcal{B}_{qj}^T + \mathcal{B}_{jq} \mathcal{B}_{qi}^T)$, where \bar{N}_q is the thermal occupancy of the bath referred to in the q th entry in \vec{r}_{in} .

Since the Hamiltonian is quadratic and the dissipators are of Lindblad form with respect to linear combinations of the annihilation and creation operators, the time evolution of the system is closed in the set of Gaussian states [46]. Furthermore, it is also known that if the dynamics of the system is stable, i.e., the real parts of all the eigenvalues of \mathcal{A} are negative [47], then the dynamics governed by the Lyapunov equation reaches a unique steady state that can be obtained by solving the equation $\mathcal{V} = 0$, which is outlined in the Supplemental Material [45].

Nonreciprocity of entanglement propagation. These results allow us to center our attention on the asymmetry of physical quantities between modes k and m . Thus, we denote $\mathcal{V}^{(k,m)}$ as the reduced covariance matrix that characterizes the subsystem consisting of nodes k and m by ignoring all other entries [48]. Accordingly, it can be further reduced to 2×2 blocks $\mathcal{V}^{(k)}$ and $\mathcal{V}^{(m)}$ that describe the covariance matrix of the respective single modes and contains the 2×2 correlation blocks $\mathcal{V}^{(km)}$. For the covariance matrix of a two-mode Gaussian state, the symplectic eigenvalues after partial transposition \tilde{v}_i of the covariance matrix such as $\mathcal{V}^{(k,m)}$ allow to judge the separability of the subsystem [48–51]. These eigenvalues can be computed with $\Delta(\mathcal{V}^{(k,m)}) = \det(\mathcal{V}^{(k)}) + \det(\mathcal{V}^{(m)}) - 2 \det(\mathcal{V}^{(km)})$ as $\tilde{v}_{\pm}^{(k,m)} = \sqrt{\Delta(\mathcal{V}^{(k,m)}) \pm \sqrt{\Delta(\mathcal{V}^{(k,m)}) - 4 \det(\mathcal{V}^{(km)})}} / \sqrt{2}$. The positive partial transpose criterion for separability of the state characterized by $\mathcal{V}^{(k,m)}$ can be stated as $\tilde{v}_{-}^{(k,m)} \geq \frac{1}{2}$, with the amount of entanglement being evaluated through the logarithmic negativity [48–51]

$$E_{\mathcal{N}}^{(k,m)} = \max\{0, -\ln(2|\tilde{v}_{-}^{(k,m)}|)\}, \quad (3)$$

which quantifies the entanglement between the two subsystems of a bipartition by being zero for a separable state and positive for an entangled state.

Our analysis proceeds with the properties of the steady-state covariance matrix $\tilde{\mathcal{V}}$, given through the solution of $\tilde{\mathcal{V}}\mathcal{A}^T + \mathcal{A}\tilde{\mathcal{V}} + \mathcal{N} = 0$ as the fixed point of the Lyapunov equation [52]. The system under consideration consists of identical oscillators $\omega_k = \omega$ and a channel of length $M = 10$ with $\gamma/\omega = 0.8$, $\gamma_{\text{out}}/\omega = 0.002$, and initial mean occupations $\tilde{N}_k = \tilde{N}_{l,l+1} = 0$. A comment is in order here regarding the counterintuitive nature of the value of γ/ω . Perfect directionality in the chain can only be achieved if the hopping rates J_l are equal to $\gamma/2$, as detailed above. Should the dissipation rate γ be too small, the hopping rate between adjacent nodes would be similarly small and the entanglement would not propagate effectively through the chain. Conversely, if γ is too large, dissipation is too strong and entanglement once again fails to propagate. The values of γ and J_l employed here strike a balance between these two scenarios and allow for the efficient propagation of entanglement. Further elaboration of the parameters is given in the Supplemental Material [45].

The behavior of the entanglement in the system is determined by the amount of squeezing of the auxiliary oscillator \hat{a}_0 , tunable through the parameter r , and its connection to the chain with regard to the coupling structure's forward direction, as characterized by J_0 and J_M . The connection of the squeezed oscillator in the forward direction of the chain is realized by $iJ_0 = j$ and $J_M = 0$, while the connection against




FIG. 2. Logarithmic negativity between the squeezed node and the node after the first nonreciprocal coupling in the forward direction (solid surface) and in the backward direction (meshed surface). The solid surface shows that connecting the squeezed oscillator to the channel in the forward direction results in nonzero logarithmic negativity indicating entanglement between the squeezed node and the nodes of the channel. The meshed surface shows that connecting the squeezed node against the forward direction results in zero logarithmic negativity irrespective of the strength of squeezing and coupling. Consequently, the squeezed oscillator and the channel are in a separable state demonstrating the effective isolation of entanglement propagation through the nonreciprocal coupling.

the forward direction of the chain requires $J_0 = 0$ and $-iJ_M = j$. We characterize the entanglement properties of the nonreciprocal coupling scheme through properties between the squeezed oscillator and the oscillator after the first nonreciprocal edge. Figure 2 illustrates the logarithmic negativity $E_{\mathcal{N}}^{(0,\pm 2)}$ between oscillator \hat{a}_0 and the oscillator two hops away, with regard to the strength of squeezing and its coupling to the channel; in other words, $E_{\mathcal{N}}^{(0,+2)} := E_{\mathcal{N}}^{(0,2)}$ quantifies the entanglement between the squeezed node and its next-to-nearest neighbor in the forward direction, and $E_{\mathcal{N}}^{(0,-2)} := E_{\mathcal{N}}^{(0,9)}$ in the reverse direction. We find that coupling the squeezed oscillator to the channel in the forward direction, depicted by the solid surface, enables nonzero values of the logarithmic negativity, and we therefore find that the corresponding nodes are entangled. Moreover, we find that both squeezing and coupling are required to achieve entanglement as squeezing creates identical excitation pairs in \hat{a}_0 that can only travel to the channel if excitations can hop to the channel's first node and increasing both parameters results in a larger logarithmic negativity. If the squeezed oscillator is coupled to the channel against the forward direction, as illustrated by the meshed surface, the resulting state between the channel and the squeezed oscillator is separable, evidenced by vanishing logarithmic negativity irrespective of the squeezing and coupling parameters. For large squeezing and hopping parameters, the system is found to be unstable, which can be seen in the edges of the surfaces resulting from the logarithmic negativity being undefined. Nevertheless, we find that entanglement of the constituents of the quantum channel




FIG. 3. Propagation depth of entanglement and mean occupation number of the final entangled node in the quantum channel for the steady state in the forward direction. (a) Maximal node in the quantum channel that is entangled with the squeezed oscillator depending on hopping and squeezing parameters. The maximal node increases with increasing the coupling of the squeezed oscillator to the quantum channel and decreases with the squeezing. For low squeezing and large hopping, entanglement passes throughout the entire system. Increasing the squeezing leads to instability of the system. (b) The mean occupation number of the maximal entangled node is related to the temperature increase generated through the entangling scheme. The mean occupation increases with larger coupling of the squeezed oscillator to the quantum channel as well as with the squeezing until the system becomes unstable. This result suggests that entangling the nodes in the system with the suggested scheme increases the temperature in the system until the system becomes unstable.

only propagates in the forward direction of the unidirectional coupling.

Influence of thermal fluctuations. Aiming to understand the amount of nodes that can be entangled through the channel and its limits, we find the maximal node M_{\max} inseparable from the squeezed node by the largest M with logarithmic negativity $E_{\mathcal{N}}^{(0,M)} > 0$. The results of this analysis are depicted in Fig. 3(a). Our results show that an increasing coherent coupling of the squeezed oscillator to the chain increases the amount of entangled nodes in the chain. We explain this increase by the improvement of correlated excitations to propagate to the chain with an enlarged hopping rate. However, we find that a stronger squeezing and therefore a larger rate of generating excitation pairs leads to fewer nodes that end up in an entangled state with the squeezed node and ultimately to an unstable system. Additionally, the single-mode covariance matrices $\mathcal{V}^{(k)}(t)$ are related to the mean occupation of each node as $\bar{N}_k(t) = [\text{Tr}(\mathcal{V}^{(k)}(t)) - 2]/4$, which enables us to characterize the thermal noise that is generated in the respective node. In the following, we consider the mean occupation $\bar{n}_k = [\text{Tr}(\tilde{\mathcal{V}}^{(k)}) - 2]/4$ using the steady state [52]. We attribute this occupation to the thermal noise, which is generated through the creation and propagation of entanglement throughout the quantum channel in our scheme. The resulting mean occupation in the maximal node inseparable from the squeezed node is depicted in Fig. 3(b). We find that the mean occupation number and therefore the temperature in the node rises with an increasing hopping rate and with an increasing squeezing parameter. Ultimately, the figure shows that the system becomes unstable when the occupation number becomes too large. In conclusion, we find that an increased squeezing, the same mechanism that generates entanglement, also increases the effective temperature in the system. This temperature increase causes instability in the system and turns out to be the limiting factor in the propagation of entanglement through the system.

Optomechanical implementation. This section aims to employ prior results [42,53] showing that mechanical degrees of freedom can act as controllable reservoirs required for our entanglement scheme. Consequently, the optomechanical system can be used as an entanglement routing channel for the optical fields. We employ the mapping worked out in Ref. [42], which shows that the nonreciprocal coupling can be achieved with dielectric nanostring mechanical resonators [17] or on-chip microwave electromechanical systems based on a lumped-element superconducting circuit with a drumhead capacitor [16,18]. The resulting parameters amount to cavities resonant at $2\pi \times 5$ GHz with damping rates $\kappa = 2\pi \times 2$ MHz, mechanical resonance frequency $\omega_m = 2\pi \times 6$ MHz, and damping rate $\gamma_m = 2\pi \times 100$ Hz. Although the value of γ , to which the values of J_l are tied, as discussed previously, is significantly higher than is the norm for optomechanical experiments, we believe that such values are not outside the realm of experimental realization for systems that are designed and optimized appropriately.

Conclusion and outlook. Our investigation outlines a general framework for nonreciprocal propagation of entanglement in composite bosonic quantum systems. Our study also shows that the creation of entanglement increases the temperature in the system, which in turn is the limiting factor for the stability of the system. Our framework can be mapped to an optomechanical realization. This mapping shows that the scheme can be realized with parameters of typical microwave optomechanical experiments. Therefore, our scheme can be tested in present-day experiments with state-of-the-art setups in the optical and the microwave domain. In the context of quantum measurements and emerging quantum technologies, our techniques and ideas will find use in the manipulation of flow of entanglement inside quantum devices for phonon-based signal processing and computation.

Acknowledgments. The authors thank S. Barzanjeh for insightful discussions and feedback. K.P. and A.X.

acknowledge financial support from the Ministry for Education, Sports, Youths, Research and Innovation of the Government of Malta through its participation in the QuantERA ERA-NET Cofund in Quantum Technologies (project MQSens) implemented

within the European Union's Horizon 2020 Programme.

Data availability. The data that support the findings of this article are not publicly available. The data are available from the authors upon reasonable request.

-
- [1] M. Mohseni, P. Read, H. Neven, S. Boixo, V. Denchev, R. Babbush, A. Fowler, V. Smelyanskiy, and J. Martinis, Commercialize quantum technologies in five years, *Nature (London)* **543**, 171 (2017).
- [2] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, *Rev. Mod. Phys.* **81**, 865 (2009).
- [3] A. Acín, I. Cirac, and M. Lewenstein, Entanglement percolation in quantum networks, *Nat. Phys.* **3**, 256 (2007).
- [4] H. J. Kimble, The quantum internet, *Nature (London)* **453**, 1023 (2008).
- [5] F. Xu, X. Ma, Q. Zhang, H. K. Lo, and J. W. Pan, Secure quantum key distribution with realistic devices, *Rev. Mod. Phys.* **92**, 025002 (2020).
- [6] S. Pirandola, U. L. Andersen, L. Banchi, M. Berta, D. Bunandar, R. Colbeck, D. Englund, T. Gehring, C. Lupo, C. Ottaviani, J. L. Pereira, M. Razavi, J. Shamsul Shaari, M. Tomamichel, V. C. Usenko, G. Vallone, P. Villoresi, and P. Wallden, Advances in quantum cryptography, *Adv. Opt. Photon.* **12**, 1012 (2020).
- [7] E. Pelucchi, G. Fagas, I. Aharonovich, D. Englund, E. Figueroa, Q. Gong, H. Hannes, J. Liu, C. Y. Lu, N. Matsuda, J. W. Pan, F. Schreck, F. Sciarrino, C. Silberhorn, J. Wang, and K. D. Jöns, The potential and global outlook of integrated photonics for quantum technologies, *Nat. Rev. Phys.* **4**, 194 (2022).
- [8] J. I. Cirac, P. Zoller, H. J. Kimble, and H. Mabuchi, Quantum state transfer and entanglement distribution among distant nodes in a quantum network, *Phys. Rev. Lett.* **78**, 3221 (1997).
- [9] J. Yin, Y. H. Li, S. K. Liao, M. Yang, Y. Cao, L. Zhang, J. G. Ren, W. Q. Cai, W. Y. Liu, S. L. Li, *et al.*, Entanglement-based secure quantum cryptography over 1,120 kilometres, *Nature (London)* **582**, 501 (2020).
- [10] A. Metelmann and A. A. Clerk, Nonreciprocal photon transmission and amplification via reservoir engineering, *Phys. Rev. X* **5**, 021025 (2015).
- [11] J. Kim, M. C. Kuzyk, K. Han, H. Wang, and G. Bahl, Nonreciprocal Brillouin scattering induced transparency, *Nat. Phys.* **11**, 275 (2015).
- [12] C. H. Dong, Z. Shen, C. L. Zou, Y. L. Zhang, W. Fu, and G. C. Guo, Brillouin-scattering-induced transparency and nonreciprocal light storage, *Nat. Commun.* **6**, 6193 (2015).
- [13] K. M. Sliwa, M. Hatridge, A. Narla, S. Shankar, L. Frunzio, R. J. Schoelkopf, and M. H. Devoret, Reconfigurable Josephson circulator/directional amplifier, *Phys. Rev. X* **5**, 041020 (2015).
- [14] F. Lecocq, L. Ranzani, G. A. Peterson, K. Cicak, R. W. Simmonds, J. D. Teufel, and J. Aumentado, Nonreciprocal microwave signal processing with a field-programmable Josephson amplifier, *Phys. Rev. Appl.* **7**, 024028 (2017).
- [15] Z. Shen, Y. L. Zhang, Y. Chen, C. L. Zou, Y. F. Xiao, X. B. Zou, F. W. Sun, G. C. Guo, and C. H. Dong, Experimental realization of optomechanically induced non-reciprocity, *Nat. Photon.* **10**, 657 (2016).
- [16] N. R. Bernier, L. D. Tóth, A. Koottandavida, M. A. Ioannou, D. Malz, A. Nunnenkamp, A. K. Feofanov, and T. J. Kippenberg, Nonreciprocal reconfigurable microwave optomechanical circuit, *Nat. Commun.* **8**, 604 (2017).
- [17] S. Barzanjeh, M. Wulf, M. Peruzzo, M. Kalaei, P. B. Dieterle, O. Painter, and J. M. Fink, Mechanical on-chip microwave circulator, *Nat. Commun.* **8**, 953 (2017).
- [18] G. A. Peterson, F. Lecocq, K. Cicak, R. W. Simmonds, J. Aumentado, and J. D. Teufel, Demonstration of efficient nonreciprocity in a microwave optomechanical circuit, *Phys. Rev. X* **7**, 031001 (2017).
- [19] L. Mercier de Lépinay, E. Damskägg, C. F. Ockeloen-Korppi, and M. A. Sillanpää, Realization of directional amplification in a microwave optomechanical device, *Phys. Rev. Appl.* **11**, 034027 (2019).
- [20] H. Xu, A. A. Clerk, and J. G. E. Harris, Nonreciprocal control and cooling of phonon modes in an optomechanical system, *Nature (London)* **568**, 65 (2019).
- [21] M. Aspelmeyer, T. J. Kippenberg, and F. Marquardt, Cavity optomechanics, *Rev. Mod. Phys.* **86**, 1391 (2014).
- [22] S. Barzanjeh, A. Xuereb, S. Gröblacher, M. Paternostro, C. A. Regal, and E. M. Weig, Optomechanics for quantum technologies, *Nat. Phys.* **18**, 15 (2022).
- [23] A. Xuereb, C. Genes, and A. Dantan, Strong coupling and long-range collective interactions in optomechanical arrays, *Phys. Rev. Lett.* **109**, 223601 (2012).
- [24] P. Piergentili, L. Catalani, M. Bawaj, S. Zippilli, N. Malossi, R. Natali, D. Vitali, and G. Di Giuseppe, Two-membrane cavity optomechanics, *New J. Phys.* **20**, 083024 (2018).
- [25] M. Schmidt, S. Kessler, V. Peano, O. Painter, and F. Marquardt, Optomechanical creation of magnetic fields for photons on a lattice, *Optica* **2**, 635 (2015).
- [26] V. Peano, C. Brendel, M. Schmidt, and F. Marquardt, Topological phases of sound and light, *Phys. Rev. X* **5**, 031011 (2015).
- [27] J. P. Mathew, J. del Pino, and E. Verhagen, Synthetic gauge fields for phonon transport in a nano-optomechanical system, *Nat. Nanotechnol.* **15**, 198 (2020).
- [28] S. Walter, and F. Marquardt, Classical dynamical gauge fields in optomechanics, *New J. Phys.* **18**, 113029 (2016).
- [29] G. Heinrich, M. Ludwig, J. Qian, B. Kubala, and F. Marquardt, Collective dynamics in optomechanical arrays, *Phys. Rev. Lett.* **107**, 043603 (2011).
- [30] R. Lauter, C. Brendel, Steven J. M. Habraken, and F. Marquardt, Pattern phase diagram for two-dimensional arrays of coupled limit-cycle oscillators, *Phys. Rev. E* **92**, 012902 (2015).
- [31] R. Lauter, A. Mitra, and F. Marquardt, From Kardar-Parisi-Zhang scaling to explosive desynchronization in arrays of limit-cycle oscillators, *Phys. Rev. E* **96**, 012220 (2017).
- [32] C. A. Holmes, C. P. Meaney, and G. J. Milburn, Synchronization of many nanomechanical resonators coupled via a common cavity field, *Phys. Rev. E* **85**, 066203 (2012).

- [33] N. Lörch, S. E. Nigg, A. Nunnenkamp, R. P. Tiwari, and C. Bruder, Quantum synchronization blockade: energy quantization hinders synchronization of identical oscillators, *Phys. Rev. Lett.* **118**, 243602 (2017).
- [34] E. Amitai, N. Lörch, A. Nunnenkamp, S. Walter, and C. Bruder, Synchronization of an optomechanical system to an external drive, *Phys. Rev. A* **95**, 053858 (2017).
- [35] M. Zhang, G.S. Wiederhecker, S. Manipatruni, A. Barnard, P. McEuen, and M. Lipson, Synchronization of micromechanical oscillators using light, *Phys. Rev. Lett.* **109**, 233906 (2012).
- [36] M. Zhang, S. Shah, J. Cardenas, and M. Lipson, Synchronization and phase noise reduction in micromechanical oscillator arrays coupled through light, *Phys. Rev. Lett.* **115**, 163902 (2015).
- [37] M. F. Colombano, G. Arregui, N. E. Capuj, A. Pitanti, J. Maire, A. Griol, B. Garrido, A. Martinez, C. M. Sotomayor-Torres, and D. Navarro-Urrios, Synchronization of optomechanical nanobeams by mechanical interaction, *Phys. Rev. Lett.* **123**, 017402 (2019).
- [38] J. Chan, T. P. M. Alegre, A. H. Safavi-Naeini, J. T. Hill, A. Krause, S. Gröblacher, M. Aspelmeyer, and O. Painter, Laser cooling of a nanomechanical oscillator into its quantum ground state, *Nature (London)* **478**, 89 (2011).
- [39] J. D. Teufel, T. Donner, D. Li, J. W. Harlow, M. S. Allman, K. Cicak, A. J. Sirois, J. D. Whittaker, K. W. Lehnert, and R. W. Simmonds, Sideband cooling of micromechanical motion to the quantum ground state, *Nature (London)* **475**, 359 (2011).
- [40] C. F. Ockeloen-Korppi, E. Damskägg, J. M. Pirkkalainen, M. Asjad, A. A. Clerk, F. Massel, M. J. Wooley, and M. A. Sillanpää, Stabilized entanglement of massive mechanical oscillators, *Nature (London)* **556**, 478 (2018).
- [41] R. Riedinger, A. Wallucks, I. Marinković, C. Löschnauer, M. Aspelmeyer, S. Hong, and S. Gröblacher, Remote quantum entanglement between two micromechanical oscillators, *Nature (London)* **556**, 473 (2018).
- [42] S. Barzanjeh, M. Aquilina, and A. Xuereb, Manipulating the flow of thermal noise in quantum devices, *Phys. Rev. Lett.* **120**, 060601 (2018).
- [43] L. Orr, S. A. Khan, N. Buchholz, S. Kotler, and A. Metelmann, High-purity entanglement of hot propagating modes using non-reciprocity, *PRX Quantum* **4**, 020344 (2023).
- [44] C. Gardiner and P. Zoller, Cascaded quantum systems, in *Quantum Noise*, 2nd ed., edited by H. Haken (Springer, New York, 2000), Chap. 12, pp. 397–414.
- [45] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/hsc8-pv2s> for the determination of the dynamic and noise matrices for the model used, the analysis of stability and physicality of the dynamical equations, and a determination of the operational parameters of the system under consideration.
- [46] T. Linowski, A. Teretenkov, and Ł. Rudnicki, Dissipative evolution of quantum Gaussian states, *Phys. Rev. A* **106**, 052206 (2022).
- [47] M. Behr, P. Benner, and J. Heiland, Solution formulas for differential Sylvester and Lyapunov equations, *Calcolo* **56**, 51 (2019).
- [48] A. Serafini, *Quantum Continuous Variables*, 1st ed. (Taylor & Francis, Boca Raton, FL, 2000).
- [49] G. Adesso, A. Serafini, and F. Illuminati, Extremal entanglement and mixedness in continuous variable systems, *Phys. Rev. A* **70**, 022318 (2004).
- [50] G. Adesso and F. Illuminati, Gaussian measures of entanglement versus negativities: Ordering of two-mode Gaussian states, *Phys. Rev. A* **72**, 032334 (2005).
- [51] A. Serafini, F. Illuminati, and S. De Siena, Symplectic invariants, entropic measures and correlations of Gaussian states, *J. Phys. B: At. Mol. Opt. Phys.* **37**, L21 (2004).
- [52] I. Pietikäinen, O. Černotik, and R. Filip, Combining Floquet and Lyapunov techniques for time-dependent problems in optomechanics and electromechanics, *New J. Phys.* **22**, 063019 (2020).
- [53] S. Barzanjeh, S. Guha, C. Weedbrook, D. Vitali, J. H. Shapiro, and S. Pirandola, Microwave quantum illumination, *Phys. Rev. Lett.* **114**, 080503 (2015).